

Immunization Training Chart

UPDATED: 5/2012

<i>Minimum interval between Doses</i>					
Vaccine	Minimum / maximum age for dose 1	Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
DTap	6 weeks- 3 months	4 weeks (age 3-5 months)	4 weeks (age 5-7 months)	6 months (age 8-19 mo.)	6 months (age 5-7yrs)
Polio	6 weeks- 3 months	4 weeks (age 3-5 months)	4 weeks age (5-19 months)	6 months (age 5-7years)	XXXXXXXXXXXX
MMR	12 - 16 months	4 weeks (age 5-7 yrs)	XXXXXXXXXXXX		
Hepatitis B	Birth- 3 months	4 weeks (age 3-5 months)	8 weeks (age 7-19 mos)	XXXXXXXXXXXX	
Rotavirus	6 weeks- 3 months	4 weeks (age 3-5months)	4 weeks (age 6-8 mos)	Not given after age 8 mo.	
Varicella	12 - 16 months	3 months If younger than 13 yrs 4 months if 13 yrs + (due at age 5-7 yrs)	XXXXXXXXXXXX		
Hepatitis A	12 - 16 months	6 months (due at 22-24 months)	XXXXXXXXXXXX		
Hib	6 weeks- 3 months	4 weeks If 1 st dose given before age 12months (4-12 months of age) 8 weeks if last dose was given at 12-14 mo. If first dose was given after 15 mo. no further doses needed	4 weeks If younger than 12 mos 8 weeks (as final dose) if older than 12 mo. & 2 nd dose given at younger than 15 mo. No further doses needed if previous dose was given at 15 mos. or older	8 weeks (as final dose) (18-60 months of age) this dose only necessary for children aged 12 -59 months who have received 3 doses before age 12 mo.	
Pneumococcal	6 weeks-3 months	4 weeks if 1 st dose administered at younger than 12 months (4-12mo age) 8 weeks (as final dose for healthy children) If 1 st dose was administered at age 12 months/ or 24- 59 months No further doses needed	4 weeks if less than 12 months (12-18 ms of age) 8 weeks (Final dose for healthy children) If current age is 12 months or older No further doses needed	8 weeks (as final dose) This dose only necessary for children age 12 thru 59 months who have received 3 doses before age 12 months—or— for high risk who have received 3 doses at any age	

1. **DTap:** The 5th dose is not necessary if the 4th dose was administered at age 4 years or older
2. **Polio:** The 4th dose is not necessary if the 3rd dose was administered at age 4 years or older and at least 6 months following the previous dose.
3. **Hep B:** Minimum age for the 3rd dose is 24 weeks and at least 16 weeks after the first dose
4. **Rotavirus:** The maximum age for the first dose is 14 weeks 6 days. Vaccination should not be initiated for infants aged 15 weeks 0 days or older. The maximum age for the final dose in the series is 8 months 0 days. If Rotarix was administered for the first and second doses, a third dose is not indicated.
5. **Pneumococcal:** Children aged 14-59 months who received a completed 4- dose or other age appropriate series of PCV7, must also receive a supplemental dose of PCV13.